INDEX TO VOLUME XII

New species in blackfaced type

AAMODT, O. S., Correlated inheritance in wheat of winter-spring habit of growth and rust resistance, 32; The inheritance of resistance to several biologic forms of Puccinia graminis tritici in a cross between Kanred and Marquis wheats, 32; see STAKMAN, E. C.

Abstracts of paper presented at the thirteenth annual meeting of the American Phytopathological Society, Toronto, Can., Dec. 28–31, 1921, 27–60; fifth annual meeting of the Pacific Division, Berkeley, Cal., Aug. 4–6, 1921, 103–8; fourth annual meeting of the Southern Division, Atlanta, Ga., Feb. 20–22, 1922, 250–2; sixth annual meeting of the Pacific Division, Salt Lake City, Utah, June 22–24, 1922, 442–448

ADAMS, J. F., Observations on frost protection and drouth spot of apple, 184–187; and MANNS, T. F., The corn ear worm and kernel rot of corn, 25–26

Adelia acuminata., host of Pseudomonas savastanoi, 106, 271–8

Air, upper, spore content of, 44

Alfalfa, girdle, experiments on, 188–190; infestations by nematode, 105

ALLEN, RUTH F. Observations on the nature of resistance to stem rust attack in certain resistant and susceptible wheats, 108; see MACKIE, W. W.

Alternaria **radicina**, cause of black rot of carrots, 157–166; cause of storage rot of carrots, 49; cause of new spot of tomato, 146–148

American Phytopathological Society, Abstracts of papers presented at the thirteenth annual meeting of, Toronto, Can., Dec. 28–31, 1921., 27–60; Abstracts of papers presented at the fifth annual meeting of the Pacific Division of, Berkeley, Cal., August 4–6, 1921, 103–108; Report of the thirteenth annual meeting of, Toronto, Can., 195–204; Abstracts of papers presented at the fourth annual meeting of the Southern Division of, Atlanta, Ga., Feb. 20–22-250–252; Report of fourth annual meeting of southern Division, 249; Abstracts of papers presented at the sixth annual meeting of the Pacific Division of, Salt Lake City, Utah, June 22–24, 1922, 442–448

Angular leaf-spot of cotton, control of, 20–22

Anthracnose, hybrid bean resistant to 47; of pea, 47; of raspberry, control of, 57

Aphis rubiphila, agent in disseminating leaf-curl and mosaic, 58; 253–64

Apple, Bellflower, susceptibility to cedar rust, 190–192; bitter pit. 489–491; blotch canker, origin of, 55; drouth spot and frost protection, 184–187; roots parasitized by Comandra pallida, 104; scab of, 54; scald, control of, 103

ARTSCHWAGER, E., Occurrence of Phloem necrosis in leafroll tubers, 193

Ascochyta gossypii Sydow, cotton blight caused by, 250

Asparagus, a new disease of, 49

Aspen, host of Fomes igniarius, 446

Aspergillus niger, infection of figs by, 442

Avena sativa, crown rust of, 4, 108; formaldehyde treatment for smut, 35, 36; resistance to stem rust attack, 108; treatment with copper carbonate dust to prevent smut, 36; host of Septoria avenae, 450–470; host of Leptosphaeria **avenaria**; 450–470

Azuki-bean (Phaseolus radiatus var. aurea) mosaic of, 295–7

Bacillus amylovorus, on Eriobotrya japonnica, 399; A new host for, morphological and cultural studies of, 520–524; summary of hosts on which reported, 519; atrosepticus, melanogenes, phytophthorus, and solanisaprus, on potato, 444

Bacterium, citriputeale, relation to environmental factors, 107; solanacearum new hosts, for, 98; tabacum, wildfire disease of tobacco, 51; tumefaciens,

effect on tobacco plants, 56; tumefaciens inoculations to cause fasciation, 265–70; sp. on Capsicum annuum, morphological and physiological studies of, 501–516

Bailey, D. L., Investigations on Puccinia helianthi Schw., 44; see Lambert, E. B.

Ball, E. D., Insects as disseminators of plant diseases, II, Systematic relations of carriers, 229–231

Banana freckle, a black spot disease, 101

Barberry, eradication campaign, progress of, 33

Barley, see Hordeum sativum

Barrett, J. T., Notes on bacterial gummosis of stone fruits, 103

Barrus, M. F., and Chupp, C. C., Yellow dwarf of potatoes, 39, 123–132

Barss, H. P., Destructive rust (Puccinia subnitens Dietel) on spinach in the Northwest, 446

Bartholomew, E. T., Acid and water content of lemon fruits at different stages of development, 107; Internal decline of lemons, 106; see also Bennett J. P.

Bastard toad flax, parasitic on apple roots, 104

Bean, hybrid resistant to anthracnose and mosaic, 47; lima see Phaseolus lunatus; parasitized by yeast, 47; soy, host of Bacterium solanacearum, 98

Becker, G. G., Three years' sweet potato certification work in Arkansas, 252

Beet leafhopper, relation to curly leaf, 105

Bellflower apple, susceptibility to cedar rust, 190–192

Bennett, J. P. and Bartholomew, E. T., Respiration of potatoes in relation to the occurrence of blackheart in storage, 443

Bensuade, M., and Keitt, G. W., A preliminary report on cross-inoculation experiments with strains of Cladosporium from stone fruits, 46

Biologic forms of Puccinia graminis, 353–361

Bisby, G. R., and Clayton, E. E., Martin, W. H., Rosa, J. T., Jr., and Stokdyk, E. A., The cooperative potato spraying project; Report for 1921, 241–248

Blackheart, of potatoes, 443

Black leg of cabbage, occurrence of, 43

Black point, of wheat, 34

Black rot, of carrot, caused by Alternaria **radicina**, 157–166; of cucumber, cause of, 43

Blair, R. J., Pulp storage in water, 60

Blight, Ascochyta, of cotton, 250; spraying for bacterial and late blight of celery, 48; needle of Pinus strobus, 58; of wheat seedlings by Fusarium culmorum, 28; twig, of elm, 47; walnut, infection and resistance to, 106

Blodgett, F. M., The relation of time and temperature to the killing of potatoes and potato mosaic virus, 40; and Perry, F. R., Additions of formalin to maintain the concentration uniform with direct steam heat in the hot formaldehyde treatment of potatoes, 39; Fernow, Karl, and Perry, F. R., Testing seed potatoes for mosaic and leaf roll II, 40

Bonar, Lee, The life history of Rosellina **caryae** causing a hickory canker and disease, 381–385

Bordeaux mixture, effect of, on size of sweet cherries, 104; spraying celery with, 48; use for crown gall, 55; use for leaf-spot of eggplant, 63; use for raspberry anthracnose, 57

Botrytis, libertiana, cause of fig disease, 107

Bottom rot disease, of lettuce, 48

Boyce, J. S., White pine blister rust in the Pacific Northwest, 448

Branstetter, B. B., treatment of seed to control root and stalk rots, 30

Braun, Harry, Effect of delayed planting on germination of seed wheat treated with formalin, 173–179

Bremia lactucae, cause of transit disease of lettuce, 48

Briggs, F. N., The toxicity of copper sulphate to the spores of Tilletia tritici

(Bjerk) Winter, 446; Stimulation by copper carbonate of wheat seedlings in the Greenhouse, 446; see MACKIE, W. W

BROOKS, CHARLES, COOLEY, J. S. and FISHER, D. F., Experiments on the use of oiled fruit wraps for the control of apple scald, 103

BROWN, J. G., and GIBSON, FREDERICK, Some observations on alfalfa girdle, 188–190

Brown rot of peaches, control of, by dusting vs. spraying, 250

Bunt, relation of soil-moisture and temperature to wheat-infection, 108; 337–52; use of chemical dusts to prevent, 108

BUTLER, O., and SMITH, T. O., On the use of the acetates of copper as fungicides, 279–89

Cabbage black leg, occurrence, of, 43

CAESAR, L., Insects as disseminators of plant diseases, III. Control problems, 231–233

Calcium chloride, injurious to Pinus strobus, 367–368

Calico, of potato, characters of, 133–139

Canker, apple blotch, origin of, 55; European, in Pacific Coast States, 105; poplar, caused by Hypoxylon primatum, 59

Capsicum annuum, Phytophthora disease of, 403–408; Bacterial spot of, 501–516

Carrot, black rot of, caused by Alternaria **radicina**, 157–166; storage rot of, 49

CARSNER, EUBANKS, Curly top of beet. Is it a mosaic disease? 105

Carya ovata, canker of, 381–385; host of Rosellinia **caryae**, 381–85; host of Dothichiza **caryae**, 381–385

Celery, damping-off of, by Sclerotinia libertiana, 16–20; mosaic, 151–4; spraying and dusting experiments on, 48; yellows, variety resistant to, 48

Cephalosporium sacchari, cause of kernel rot of corn, 25

Cercospora **melongenae**, leafspot disease of eggplant, 61–5

Cereals, present status of pathology of, in U. S., 108; Septoria diseases of, 44; take-all disease of, 66–88

Chemical dusts, use to prevent bunt, 108

Chemical injury, to various hosts, 364

Cherry, effect of alkaline spray on size of, 104

Chionanthus virginica host of Pseudomonas savastanoi, 106; 271–8

CHRISTOPHER, W. N., see STAKMAN, E. C.

CHUPP, C. C., see BARRUS, M. F.

Citrus blast, relation to climatic conditions, 107; plants, susceptibility to scab, 57

Cladosporium citri, susceptibility of citrus plants to, 57; cross-inoculation experiments with, 46

CLAYTON, E. E., Diplodia zeae as an ear and root parasite of corn, 29; see BISBY, G. R. M.

Clover, host of nematode, 105

Cocoanut bud rot in the Philippines, 46; see also Cocos nucifera

Cocos nucifera, host of Thielaviopsis paradoxa, 398

Colletotrichum lindemuthianum, bean resistant to, 47; spore production in, 390–393; pisi on garden peas, 47

Comandra pallida, parasitic on apple roots, 104

Coniferae, resin canals of, 59

COOK, M. T., The dissemination of peach yellows and little peach, 140–142; A new disease of asparagus, 49; and POOLE, R. F., Stem rot disease of sweet potatoes in New Jersey, 51

COOLEY, J. S., see BROOKS, CHARLES.

COONS, G. H., and NELSON, RAY, First progress report on yellows-resistant golden self-blanching celery, 48

Copper acetates, as fungicides, 279–89

Copper carbonate dust for smut in wheat and oats, 36

Corn, see Zea mays

Corn ear worm, cause of kernel rot, 25

Corticium vagum, on potato, 444

Cotton, Ascochyta blight of, 250; control of angular leafspot of, 20–2, 50; Texas root-rot of, 250; wilt, a seed borne disease, 50

Cosmos, new host of Bacterium solanacearum, 98
Crawford, R. F., see Elliott, J. A.
Crimea, notes on fruit diseases in, 533–535
Cronartial stages, new, of two important pine cone rusts, 109–122
Cronartium **conigenum,** description and distribution, 116–122
Cronartium, ribicola, conditions necessary for infection of Pinus strobus, 148–50; effect on Ribes, 45; notes on, 46; potential sporidia production per unit, 298–9; viability of telia of, in winter, 221–4; occurrence of, in Europe, 302–4; distribution in Pacific Northwest, 448; **strobilinum** description and distribution, 109–116
Crop protection institute, fellowships offered, 156
Crown gall, infection and control of, on apple grafts, 55; studies of, 55; cause of fasciation and prolepsis, 265–70
Crown-rust of oats, data on germination of spores, 108; infection capabilities of, 4; miscellaneous studies with, 108
Cucumber, black rot, caused by Mycosphaerella citrullina, 43; mosaic of 42; relative susceptibility of European and American varieties to bacterial wilt, 143–6
Cucurbit mosaic, experiments with, 42
Cultures, plate, printing of, 53; tube, method of photographing, 399; of fungi, a new source of, 24
Curculio of peaches, control by dusting vs. spraying, 250
Curly top of beet, 105
Curran, G. C., see Stakman, E. C.
Currant, effect of Cronartium ribicola upon, 45

Dahlia, new host of Bacterium solanacearum, 98
Damping-off, as cause of root-rot in pine seedlings, 213–20; of celery by Sclerotinia libertiana, 16–20
Davis, W. H. Staining germinating spores, 492–494
Decay, of vegetables and fruits by different species of Rhizopus, 205–212
Detmers, Freda, A preliminary report on a serious twig blight of American elm, 47
Dickson, B. T., Further studies on mosaic —I, 42; and McRostie, G. P., Further studies on mosaic—II, 42
Die-back of loganberry, 104
Diplodia cacaoicola on Hevea brasiliensis, 388; zeae, ear and root parasite of corn, 29; in Iowa, 29
Doolittle, S. P., Comparative susceptibility of European and American varieties of cucumbers to bacterial wilt, 143–146; and Walker, M. N., Notes on cucurbit mosaic, 42
Dothichiza **caryae**, on Carya ovata, 381–385; populea, on populus, 425–426
Douglas, Bruce, A new Alternaria spot of tomatoes in California, 146–148
Drechsler, Charles, A new leaf spot of Kentucky Blue Grass caused by an undescribed species of Helminthosporium, 35; Net blotch of meadow fescue caused by an undescribed species of Helminthosporium, 35; see Meier, F. C.
Drouth spot, of apple and frost protection, 184–7
Dufrenoy, Jean, The occurrence of Cronartium ribicola in Europe, 302–4
Durrell, L. W., Diplodia of corn in Iowa, 29
Dusting, of potato, results of cooperative project, 241–8
Dye, H. W., The bottom-rot disease of western New York lettuce; 48; and Newhall, A. G., Spraying and dusting for the bacterial and late blights of celery in western New York, 48

Eddy, E. D., see Meier, F. C.
Egg plant, Cercospora leaf spot of, 61–5
Elliott, J. A., Cotton wilt, a seed-borne disease, 50; The Ascochyta blight of cotton, 250; and Crawford, R. F. The spread of tomato wilt by infected seed, 428–434
Elm, twig blight of, 47
Empoasca mali, injury of potatoes by, 37; cause of potato tipburn, 181–4

Erodium moschatum, host of Synchytrium papillatum, 442
EVANS, N. S., "Black point" of wheat, 34
Exoascus sp. on Prunus subcordata, 443
EYER, J. R., Notes on the etiology and specificity of the potato tip burn produced by Empoasca mali Le Baron, 181–184
EZEKIEL, W. N., Photographing tube cultures, 399

Fasciation due to crown gall, 265–70
FAULL, J. H., Records for four years on the needle blight of Pinus Strobus, 58
FAWCETT, H. S., A phomophsis of citrus in California, 107; The relation of citrus blast to certain environmental factors, 107; A new Phomopsis of citrus in California, 419–424; The relation of temperature and other factors to the etiology of plant diseases, 448
FERNOW, KARL, see BLODGETT, F. M.
Fertilizers, their effect on stem rust of wheat, 31
Fescue, meadow, net blotch of, 35
Festuca elatior, net blotch of, caused by undescribed Helminthosporium sp., 35
Fig diseases, 107; 442
Fire blight, see Bacillus amylovorus
FISHER, D. F., An outbreak of powdery mildew (Podosphaera leucotricha) on pears, 103; Effect of alkaline sprays on the size of sweet cherries, 104; The parasitism of bastard toad flax (Comandra pallida A. DC.), 104; see BROOKS, CHARLES.
FITZPATRICK, H. M., THOMAS, H. E., and KIRBY, R. S., Ophiobolus cariceti (Berk & Br.) Sacc., cause of take-all of wheat, 27
Flax wilt, 409–413
FOLSOM, DONALD and SCHULTZ, E. S., Potato tipburn in northeastern Maine, 36; see SCHULTZ, E. S.
Fomes igniarius, on Aspen, 446
Foot-rot disease of wheat in Kansas, 27
Formaldehyde, hot, treatment of potatoes, 39; treatment, of oats and wheat, 36; for prevention of oat smut, 35; for root rot, 49; for onion smut, 155; of potato unsuccessful under irrigation, 39; of seed-wheat, affected by delayed planting 173–9; use in control of Sclerotinia libertiana, 19–20
Fraxinus americana, host of Torula ligniperda, 369–374; floribunda, host of Pseudomonas savastanoi, 106; 271–278; velutina, host of Pseudomonas savastanoi, 106; 271–278
FREDERICH, W. J., see PELTIER, G. L.
FROMME, F. D., Susceptibility of apple root-stocks to black root rot, 54
Frost protection and drouth spot of apples, 184–7
FULTON, H. R., Occurrence of Thielaviopsis paradoxa on the cocoanut palm in Florida, 398–399
Fungi, damping-off, cause of root rot in pine seedlings, 213–20; new source of pure cultures of, 24; Wood-inhabiting, staining method for 440–441
Fungicides, use of the copper acetates, 279–89
Fungus cultures, a new source of supply, 24
Fusarium, culmorum, cause of wheat seedling blight, 28; infected seed corn tested in improved rag-doll germinator box, 30; lini, 409–413; lycopersici, spread of, 428–34; 30; moniliforme, cause of kernel rot of corn, 25; of corn, 251; on wheat and corn, 45; sp., cause of gladiolus rot, 53; cause of new disease of asparagus, 49; cause of stem rot of sweet potatoes, 51; variety of celery resistant to "yellows" caused by, 48; spp. causing wheat scab and relation to hydrogen-ion concentration, 290–4; sp. on spinach, 447

GARDNER, M. W., Origin of apple-blotch cankers, 55; and KENDRICK, J. B., Overwintering of tomato mosaic, 41; Insects as disseminators of plant diseases, IV. Urgent problems of the future, 233–239
GARDER, W. A., Notes on the physiology of the sweet potato, 251
Germany, plant protection service in, 23–4
Germinator box, improved rag-doll type, 30

GIBSON, FREDERICK, see BROWN, J. G.
GILBERT, A. H., The correlation of foliage-degeneration diseases of the Irish potato with variations of the tuber and sprout, 40
Gladiolus, Fusarium rot of, 53
Glume blotch, of wheat, 537–558
Gloeosporium alborubrum, on Hevea brasiliensis, 388
GODFREY, G. H., The stem and bulb infesting nematode in America, 52
Grasses, take-all disease of, 27
Grapefruit, notes on two cars from Isle of Pines, 106; host of Phomopsis **caribaea** and Diplodia natalensis, 414–417
GREGORY, C. T., The relation of rain to the formaldehyde treatment of onion smut, 155
Gum, red, host of Torula ligniperda, 370
Gummosis, bacterial, of stone fruits, 103
Gymnosporangium sabinae, reported from Crimea, 534

HAENSELER, C. M., A new peach wilt disease, 56
HAHN, G. G., see HEDGCOCK, G. G.
HARTER, L. L., Field and storage diseases of the sweet potato and their control, 251; and WEIMER, J. L. The decay of various vegetables and fruits by different species of Rhizopus, 50; Decay of various vegetables and fruits by different species of Rhizopus, 205–212; see LAURITZEN, J. I.
HASKELL, R. J., Relation of plant pathology to the newly organized Science Service, 301–2
Haskell's method, (dry formaldehyde treatment) for oat smut, 35
Heat, effect of, on wood-destroying fungi, 59
HEDGCOCK, G. G., and HAHN, G. G., Two important pine cone rusts and their new cronartial stages, Part I, Cronartium strobilinum (Arthur) Hedge and Hahn, comb. nov., 110–116; and HUNT, N. R., Part II, Cronartium conigenum (Pat.) Hedge. and Hunt, comb. nov., 116–122
Helminthosporium, disease of wheat, 28; 447; gramineum, on Hordeum sativum, 446; heveae on rubber trees, 60; oryzae, causing sesame spot disease of rice, 34; sativum, optimum temperature for, 447; sp., causing black point of wheat, 34; sp., causing seedling blight of corn affected by soil-temperature, 30; sp. undescribed, causing leaf spot of Kentucky bluegrass, 35; sp., undescribed, causing net blotch of meadow fescue, 35
HEMMI, TAKEWO, On the occurrence of Mycosphaerella wilt of muskmellons in Japan, 394–397
HENRY, A. W., see STAKMAN, E. C.
HERBERT, D. A., Bitter pit in apples: The crushed cell theory, 489–491
Hevea brasiliensis, host of Helminthosporium heveae, 60; Lightning injury to, 386–389; host of Diplodia cacaoicola, 388; host of Gloeosporium alborubrum, 388
Hickory, see Carya
HIGGINS, B. B., The bacterial spot of pepper, 501–516
HOCKEY, J. F., see RANKIN, W. H.
HOERNER, G. R., Data relative to the germination of aeciospores, urediniospores, and teliospores of Puccinia coronata Cda., 108; Infection capabilities of crown rust of oats, 4–15, 33; Miscellaneous studies with crown rust of oats, 108
HOPKINS, E. F., Varietal susceptibility of the yellow Bellflower apple to cedar rust, 190–192; The effect of lactic acid on spore production by Colletotrichum lindemuthianum, 390–393
Hordeum sativum, host of Helminthosporium spp., 446–447
HORNE, W. T., A Phomopsis from the Isle of Pines, 105; Some notes on two cars of grapefruit from the Isle of Pines, 106; A Phomopsis in grapefruit from the Isle of Pines, W. I., with notes on Diplodia natalensis, 414–418
Horse radish, root rot of, 49
HOWARD, N. O., see SNELL, WALTER H.
HOWITT, J. E., and STONE, R. E., Experiments with Haskell's method or the so-called dry formaldehyde treatment

for the prevention of oat smut, 35; see STONE, R. E.
HUBERT, E. E., A staining method for hyphae of wood inhabiting fungi, 440–441
Humidity, a factor in growth of Diplodia zeae, 29
HUMPHREY, H. B., The present status of cereal pathology in the United States, 108
HUNGERFORD, C. W., Leaf roll, mosaic, and certain other related diseases in Idaho, 133; The relation of soil moisture and soil temperature to bunt infection in wheat, 108; 337–52; Hosts for Puccinia glumarum E. & H. in the United States, 444; A fusarium blight of spinach, 447; The relation of temperature and other factors to the etiology of plant diseases, 448; see RAEDER, J. M.
HUNT, N. R., and O'DONNELL, F. G., Soil temperatures obtained under a steam pan, 53; see HEDGCOCK, G. G.
HURSH, C. R., The relation of temperature and hydrogen-ion concentration to urediniospore germination of biologic forms of stem rust, of wheat, 353–361
Hydathodes, relation to potato tipburn, 305–33
Hydrogen-ion concentration, relation to the growth of the wheat scab organism, 290–4; relation to spore germination, 353–361
Hypoxylon primatum, cause of poplar canker, 59

Infection of wheat seedlings by Helminthosporium at high temperature, 28
Inheritance of resistance, to black stem rust in crosses between varieties of common wheat, 31, 32
Insects as disseminators of plant diseases, 225–39; 1. Results of past investigations, 2. Systematic relations of carriers, 3. Control problems, 4. Urgent problems of the future
Ipomoea batatas, decay of, by Rhizopus sp., 51; diseases of, and their control, 251; notes on physiology of, 251; stem rot disease, 51; three years' certification work in Arkansas, 252
Isaria **clonostachoides**, description, characters, morphology, effect on tomato, 167–72
IYENGAR, M. O. P., and NARASIMHAN, M. J. A new species of Schizonella, 435–438

Jasminum primulinum host of Pseudomonas savastanoi, 106; 271–8
Jelly-end rot, of potato-tuber by Rhizoctonia solani, 334–6
JENNISON, H. M., Potato blackleg with special reference to the etiological agent, 444
JOHNSON, JAMES, Experimental evidence relating to the nature of the mosaic virus 54; Non-parasitic leaf spots of tobacco, 52; The relation of air temperature to the mosaic disease of potatoes and other plants, 438–440
JONES, E. S., Relation of temperature, soil moisture, and oxygen to the germination of the spores of Ustilago avenae, 45
JONES, F. R., see VAUGHAN, R. E.
JONES, L. K., A preliminary report on the control of raspberry anthracnose, 57
JONES, L. R., Whetzel resigns, 499–500; and TISDALE, W. B., The influence of soil temperature upon the development of flax wilt, 409–413
Juniperus monosperma, mistletoe on, 535–536

KEITT, G. W., Third progress report on apple scab and its control in Wisconsin, 54; see BENSAUDE, M.
KEMPTON, F. E., Progress of the barberry eradication campaign, 33
KENDRICK, J. B., see GARDNER, M. W.
Kernel rot of corn, 25
KIRBY, R. S., The improved rag-doll germinator box, 30; The take-all disease of cereals and grasses, 66–68, 27; see FRITZPATRICK, H. M.

Lactic acid, effect of, spore on production in Colletotrichum lindemuthianum, 390–393

LAMBERT, E. B., and BAILEY, D. L., Result of treating seed of spring wheat and oats with copper carbonate dust to prevent smut, 36
LARUE, C. D., Helminthosporium heveae Petch, in Sumatra, 60; Lightning injury to Hevea brasiliensis, 386–389
LAURITZEN, J. I., and HARTER, L. L., The species of Rhizopus responsible for the decay of sweet potatoes under storage conditions, 51
LEACH, J. G., Leaf hopper injury of potatoes, 37
Leaf, curl of raspberry, 58; 253–64; hopper injury of potatoes, 37; hopper the questioned cause of potato tipburn, 305–33; roll, testing seed potatoes for, II, 40; roll of potato in Idaho, 133–9; roll tubers, occurrence of phloem necrosis in, 193; rust of rye, 33; scorch, of strawberry, 375–380
Leaf-spot, angular, of cotton, 20–2, 50; non-parasitic, of tobacco, 52; of eggplant, 61–5; of Kentucky bluegrass, 35; of tobacco, 99
Leak, a transit disease of potatoes, 38
LEE, H. A., Banana freckle in the Philippine Islands, 101–102
LEIGHTY, C. E., see MAINS, E. B.
LEONIAN, L. H., Stem and fruit blight of peppers caused by Phytophthora **capsici**, 403–408.
Lemon, acid and water content of, 107; physiological diseases of, 106; host of Phomopsis **californica**, 419–423
Leptosphaeria **avenaria**, ascigerous stage of Septoria avenae, 454; physiological and taxanomic study of, 445–470; inoculation experiments on various hosts, 463–467
Lettuce, bottom-rot disease of, 48, a transit disease of, 48
LEVINE, MICHAEL, Studies on plant cancers, IV—The effect of inoculating various quantities of different dilutions of Bacterium tumefaciens into the tobacco plant, 56
Lightning injury, to Hevea brasiliensis, 386–389
Ligustrum ovalifolium, host of Pseudomonas savastanoi, 106; 271–8
Lime-sulfur, effect of, on size of sweet cherries, 104; used for raspberry anthracnose, 57
LINK, G. K. K., downy mildew; A transit disease of lettuce, 48; Leak, a serious transit disease of potatoes, 38; Transit diseases; An important factor in the cost of fresh vegetables, 105
Liriodendron tulipifera, host of Torula ligniperda, 369–374
Lister Institute, a new source of pure cultures of fungi, 24
Loganberry, die-back of, 104
LONG, W. H., Mistletoe and smelter smoke, 535–536
LUDWIG, C. A., The control of angular leaf spot of cotton, 20, 50
Lupine, (Lupinus suksdorfii) parasitized by Comandra pallida, 104
LUTMAN, B. F., The relation of the water pores and stomata of the potato leaf to the early stages and advance of tipburn, 305–33

MACINNES, JEAN, The growth of the wheat scab organism in relation to hydrogen-ion concentration, 290–4
MACMILLAN, H. G., Potato seed treatments in western states, 39; Influence of the meteorological factors on potato disease and production in Colorado, 445
MCCLINTOCK, J. A., The control of Root-knot, A progress report, 250
MCCURRY, J. B., see RANKIN, W. H.
MCKAY, M. B., Occurrence of Tylenchus dispaci on alfalfa in Oregon, 105; Thielavia basicola on watermelon in Oregon, 445; Distribution of Tylenchus dipsaci on wild strawberry in Oregon—Preliminary report, 445
MCKINNEY, H. H. and MELCHERS, L. E., Foot-rot disease of wheat in Kansas, 27
MCMURPHY, JAMES, Notes on Synchytrium, 442
MCROSTIE, G. P., see DICKSON, B. T.
MACKIE, W. W., and ALLEN, R. F., Resistance in oats to stem rust attack, 108

and BRIGGS, F. N., The use of chemicla dust to prevent bunt, 108
Macrosporium parasiticum, pathogenicity of, 50
MAINS, E. B., and LEIGHTY, C. E., Rye resistant to leaf rust, Puccinia dispersa, 33
MANEVAL, W. E., Germination of Teliospores of rusts at Colubmia, Missouri, 471–488
MANEY, T. J., see MELHUS, I. E.
MANNS, T. F., see ADAMS, J. F.
Marsonia potentillae, on strawberry, 377
MARTIN, J. P., Variation in color of pear blight exudate, 399–400
MARTIN, W. H., Further experiments with inoculated and uninoculated sulfur for the control of potato scab, 38; see BISBY, G. R.
MASSEY, L. M., Fusarium rot of gladiolus, 53
MATSUMOTO, TAKASHI, Some experiments with azuki-bean mosaic, 295–7
MEIER, F. C., and DRECHSLER, CHARLES, EDDY, E. D., Black rot of carrots caused by Alternaria **radicina**, 157–167; cucumber black rot caused by Mycosphaerella citrullina, 43; Storage rot of carrots caused by a new species of Alternaria, 49
MEINECKE, E. P., Pathology of quaking aspen in Utah in relation to regulation, 446
MELCHERS, L. E., and PARKER, J. H., Inheritance of resistance to black stem rust in crosses between varieties of common wheat (Triticum vulgare), 31; see MCKINNEY, H. H.
MELHUS, I. E., Mosaic studies, 42; and MANEY, .T J., Studies on the infection and control of crown gall on apple grafts, 55
Mercuric chloride, for angular leaf spot of cotton, 20–22; for root and stalk rot, 30; treatment for root rot, 49
Mildew, downy, cause of transit disease in lettuce, 48; powdery, outbreak on pears, 103
Moisture, relation to spore germination, 472–473
Mollisia earliana, perfect stage of Marsonia potentillae, 377
MORSTATT, H., The Biologische Reichanstalt at Berlin-Dahlem, and the plant protection service in Germany, 23
Mosaic, bean resistant to, 47; cucurbit, experiments on, 42; and leafroll, testing seed potatoes for—II, 40; of celery, 151–4; of clover, studies on, 42; of corn, 250, 252; of potato in Idaho, 133–9; of raspberry, 58; 253–64; of tomato, overwintering of, 41; diseases, in relation to temperature, 438–440; transmission by Physalis longifolia, 42; virus, nature of, 52; virus of potato, relation of time and temperature to killing of potato and virus, 40; of azuki-bean, 295–7
Moss, E. H., Observations on two poplar cankers in Ontario, 425–427
Musa sapientum, host of banana freckle disease, 101
Muskmelon, host of Mycosphaerella citrullina in Japan, 394–397
Mycosphaerella, citrullina, cause of cucumber black rot, 43; on muskmelons in Japan, 394–397

NARASIMHAN, M. J., see IYENGAR, M. O. P.
Nectria galligena, canker on apple, 105; and coccinea, morphological differences, 442; ditissima, reported from Crimea, 534
NELSON, RAY, see COONS, G. H.
Nematodes, a stem and bulb-infesting species, 52; causing root-knot, control of, 250; on alfalfa, 105; on strawberry, 445
Nematospora parasitic on lima bean, 47; **phaseoli**, on lima bean, 252–532
Net blotch, of meadow fescue, 35
NEWHALL, A. G., see DYE, H. W.

Oats, see Avena sativa
OCFEMIA, G. O., The sesame spot disease of rice, 34
O'DONNELL, F. G., Printing plate cultures, 53; see HUNT, N. REX
O'KANE, W. C., Crop Protection Institute fellowships, 156

Olive knot organism, pathogenicity on related hosts, 106; 271–8
Onion, smut, formaldehyde treatment and relation to rain, 155
Ophiobolus cariceti, cause of take-all of wheat, 27; cause of take-all of grasses and cereals, 27; cause of take-all of cereals, 66–68
Orton, C. R., The dissemination of plant diseases by seed, 54; see Weiss, Freeman
Osmanthus aquifolia host of Pseudomonas savastanoi, 106; 271–8
Owens, C. E., see Zeller, S. M.
Ozonium omnivorum, life history as affecting cotton, 250

Pacific Division, American Phytopathological Society, meeting of, see American Phytopathological Society
Parker, J. H., see Melchers, L. E.
Paxton, G. E., Studies on Helminthosporium species found on cultivated barley in California, 446
Pea, Colletotrichum pisi on, 47
Peach, decay due to Rhizopus spp., 205–12; control of curculio, brown rot, and scab, by dusting vs. spraying, 250; little peach, dissemination of, 140–2; new wilt disease of, 56; roots of, parasitized by Comandra pallida, 104; yellows; dissemination of, 140–2
Peacock, N. D., The less common spray materials, 251
Pears injured by powdery mildew, 103
Pear blight, note on color of exudate, 399
Peltier, G. L., and Frederich, W. J., Relative susceptibility of citrus plants to Cladosporium citri Massee, 57; Weather and its relation to citrus scab epidemics in Alabama, 57
Pennington, L. H., The effect of Cronartium ribicola upon Ribes, 45
Peppers, see Capsicum annuum
Perry, F. R., see Blodgett, F. M.
Personals, 156; 194
Phaseolus lunatus, yeast spot of, 525–532; host of Nematospora phaseoli, 525–532
Philippines, cocoanut bud rot in, 46
Phillips, E. H., Some fig diseases, 107; see Smith, E. H.
Phloem necrosis, in leafroll tubers and stolons, 193
Phomopsis of citrus in California, 107; a species from Isle of Pines, 105; **caribaea** on grapefruit, 147; **californica,** on lemons 419–423; temperature relations of, compared with P. citri, 422
Phorodendron juniperinum, smelter injury to, 535–536
Photographing tube cultures, 399
Phragmidium potentillae-canadensis, teliospore germination in, 478
Phyllosticta nicotiana, a leafspot disease of tobacco, 99
Physalis, a factor in transmission of tomato mosaic, 41; longifolia, agent in mosaic transmission, 42
Phytopathological Society, American, see American Phytopathological Society
Phytophthora, cause of bud rot, 46; **capsici** cause of blight of peppers, 403–408
Pine, cone rusts, new cronartial stages of two, 109–122; seedlings, root rot of, 213–20; white, see Pinus strobus; western yellow, Polyporus volvatus on, 494–496
Pinus strobus, chemical injury to, 59; 362–368; conditions necessary for infection by Cronartium ribicola, 148–150; needle blight of, 58
Plant, cancers, effect of inoculating tobacco with Bacterium tumefaciens, 56; diseases, dissemination by seed, 54; diseases, insects as disseminators of, 225–39
Plant protection service, in Germany, 23
Plum, decay due to Rhizopus spp., 205–12
Poa pratensis, new leaf spot of, 35
Podosphaera leucotricha, on pears, 103
Polyporus volvatus, on western yellow pine, 494–496
Poole, R. F., Bacterial rot of horseradish in New Jersey, 49; Celery mosaic, 151–154; Some recent investigations on the control of Sclerotinia libertiana in the greenhouse on the

muck farms of Bergen County, New Jersey, 16–20; see COOK, M. T.
Populus sp. hosts of Dothichiza populea and Cytospora chrysosperma, 425–427
Poronidulus conchifer, cause of twig blight, 47
PORTE, W. S., see PRITCHARD, F. J.
Potato, Blackleg, 444; calico of, in Idaho, 133–9; cooperative spraying project, 241–8; correlation of foliage-degeneration diseases with variations of tuber and sprout, 40; diseases in relation to weather conditions, 445; host of Corticium vagum, 444; hot-formaldehyde treatment of, 39; leaf hopper, injury of, 37; leaf-roll of, in Idaho, 133–9; leak, a transit disease of, 38; mosaic of, in Idaho, 133–9; in relation to temperature, 438–440; mosaic virus of, 40; russet dwarf of, 133–9; respiration of, in relation to blackheart, 443; scab treatment with sulphur 38; seed treatment in western states, 39; skin spot of,—a stage of powdery scab, 37; streak, transmission of, 41; testing for mosaic and leafroll (II), 40; tipburn, caused by Empoasca mali Le Baron, 181–4; tipburn in northeastern Maine, 36; tipburn, its relation to water pores and stomata, 305–33; varieties immune to wart disease, 38; wart disease investigations, 38; yellow dwarf, of 39, 123–32
Potato, sweet, decay of, by Rhizopus spp., 51; field and storage diseases of, 251; notes on physiology of, 251; stem rot disease of, 51; three years of certification work in Arkansas, 252
Potato-tuber, jelly-end rot of, 334–6
POVAH, A. H. W., Hypoxylon poplar canker, 59
Pre-soaking, in formalin-treatment of seed wheat, 173–9
Printing plate cultures, 53;
PRITCHARD, F. J., and PORTE, W. S., Isaria rot of tomato fruits, 167–172
Prolepsis, due to crown gall, 265–70
Prunus subcordata, host of new species of Exoascus, 443; triloba var. plèna, host of Bacillus amylovorus, 519–524
Pseudomonas, Cerasus on stone fruits, 103; juglandis, infection of and resistance to, 106; savastanoi, pathogenicity on hosts related to the olive, 106; 271–8; dissolvens, on stalks of Zea mays, 497–499
Puccinia, coronata, data on germination of spores, 108; coronata, infection capabilities of, 4; dispersa, resistance to, by rye, 33; graminis tritici, inheritance of resistance to, in wheat 31, 32; uredinio-spore germination in, 353–361; helianthi, 44; sorghi, 89–97; subnitens, on spinach, 446; teliospore germination in ten species of, 471–488

RAEDER, J. M., A Helminthosporium root rot of wheat in Idaho, 447; and HUNGERFORD, C. W., The effect of presprinkling with water upon the efficiency of certain potato seed treatments for the control of Rhizoctonia, 447
Rainfall, relation of, to late light of sugar-beet, 443
RAND, F. V., Insects as disseminators of plant diseases, I. Results of past investigations, 223–228
RANKIN, W. H., HOCKEY, J. F., and MCCURRY, J. B., Leaf curl and mosaic of the cultivated red raspberry, 58; and HOCKEY, J. F., Mosaic and leaf curl (yellows) of the cultivated red raspberry, 253–264
Raspberry, anthracnose, control of, 57; leaf curl of, 58; mosaic of, 58; mosaic and leafcurl (yellows) of the cultivated red raspberry, 253–64
RATHBUN, A. E., Root rot of pine seedlings 213–220
REDDICK, DONALD, A hybrid bean resistant to anthracnose and to mosaic, 47
REINKING, O. A., Cocoanut bud rot in the Philippines, 46
Rhizoctenia, cause of bottom rot disease, 48; R. solani, cause of potato-tuber rot, 334–6
Resin canals, pathological occurrence and development of, 59
Resistance to black stem rust inherited in crosses of wheat, 31, 32; to rust and

winter-spring habit of growth in inheritance, 32
Rhizoctonia, on potato seed, control of, 447
Rhizopus, decay caused by different species of, 50; 205–12; tritici and nigricans, cause of decay in sweet potatoes, 51
RHOADS, A. S., The occurrence and development of pathological resin canals in the Coniferae, 59
Ribes, effect of Cronartium ribicola on, 45; nigrum, relation to white pine blister rust, 221–4; nigrum, sporidia production in Cronartium ribicola, 298–9
Rice, diseases of, 108; sesame spot disease of, 34
RICHARDS, B. L., Relation of rainfall to the late blight or Phoma rot of the Sugar-beet, 443; Corticium vagum as a factor in potato production, 444
RIKER, A. J., Studies of crown gall, 55
Root knot, control of, 250
Root rot, black, susceptibility of apple root-stocks, to, 54; of corn, control of, by disinfection, 30; of horse radish, 49; of pine seedlings, 213–20; Texas, of cotton, 250
ROSA, J. T. JR., see BISBY, G. R.
ROSE, J. P., A seedling blight caused by Fusarium culmorum var. leteius Sher, 28
Rosellinia **caryae** on Carya ovata, 381–385
ROSEN, H. R., Corn mosaic in Arkansas, 250; "Mosaic" disease of corn in Arkansas, 252; The bacterial pathogen of corn stalk rot, 497–499
ROSTRUP, FREDERICK GEORG EMIL, biographical sketch and portrait, 1–3
Rot, Isaria rot of tomatoes, 167; jelly-end rot of potato-tuber by Rhizoctonia solani, 334–6
Russet dwarf, of potato in Idaho, character, transmission of, 133–139
Rust, black stem rust, inherited resistance to, in wheat, 31–32; cedar, urediniospore germination in 353–361; on yellow Bellflower apple, 190–2; corn, studies on, 89–97; infection affecting water requirement of wheat, 31; leaf, resistance to, by rye, 33; pine cone, new cronartial stages of two, 109–22; resistance in wheat and winter-spring habit of growth, 32; of sunflower, 44; germination of teliospores, in 471–488
Rye, resistant to leaf rust, Puccinia dispersa, 33

Sage brush (Artemesia tridentata), parasitized by Comandra pallida, 104
Scab, citrus, 57; of apple, 54; of peaches, control by dusting vs. spraying, 250; of potato, treatment with sulfur, 38; powdery, and potato "skin spot," 37
Scald, apple, method of control, 103
Schizonella **colemani**, on Vitis quadrangularis, 435–438
SCHMITZ, HENRY, Note concerning the decay of western yellow pine slash caused by Polyporus volvatus Peck, 494–497
SCHULTZ, E. S., and FOLSOM, DONALD, Transmission of potato streak, 41; see FOLSOM, DONALD
Science Service and Plant pathology, 301–2
Sclerotinia libertiana, cause of damping-off of celery, control of, 16–20; cause of fig disease, 107; spp. reported from Crimea, 533
Seed, disinfection to control root and stalk rot, 30; spread of plant diseases by, 54
Seedling blight, of corn by Helminthosporium sp. and its relation to soil-temperature, 30; of wheat by Fusarium culmorum, 28
Seedling infection of wheat by Helminthosporium greatest at high temperatures, 28
Septoria, diseases, of cereals and grasses, 44; of cereals 449–470; of wheat, 537–584; nodorum, on wheat, 537–558; inoculation experiments on numerous hosts, 550–553; tritici, on wheat, 558–584; inoculation experiments on numerous hosts, 573–575
Sesame spot of rice, 34
SEVERIN, H. H. P., Minimum incubation periods of causative agent of curly

leaf in beet leafhopper (Eutettix tenella Baker) and sugar beet, 105
SHAPOVALOV, MICHAEL, Rhizoctonia solani as a potato-tuber rot fungus, 334-6; "Skin spot," a stage of powdery scab, 37
SHEAR, C. L., F. G. E. Rostrup, 1–3; A new source of supply of pure cultures of fungi, 24
SHERBAKOFF, C. D., Fusaria of wheat and corn, 45; Fusaria of corn, 251
Shingles, asphalt, destroyed by a fungus, 497
SIGGERS, P. V., Torula ligniperda (Willk.) Sacc. a hyphomycete occurring in wood tissue, 369–374
Skin spot, of potato, a stage of powdery scab, 37
SLAGG, C. M., Preliminary report on a study of the wildfire disease of tobacco, 51
SMITH, C. O., Pathogenicity of the olive knot organism on hosts related to the olive, 106; 271–8; Some studies relating to infection of and resistance to walnut blight, Pseudomonas juglandis, 106
SMITH, E. F., Fasciation and prolepsis due to crown gall, 265–70
SMITH, E. H., and PHILLIPS, E. H., Fig smut studies, 442
SMITH, T. O., see BUTLER, O.
Smut, control of, with copper carbonate dust, 36; of oats, factors in spore germination, 45; onion, formaldehyde treatment less effective in rainy periods, 155
SNAPP, O. I., Dusting vs. spraying for the control of the curculio, brown rot, and scab of peaches, 250
SNELL, W. H., The effect of heat upon the mycelium of certain structural timber destroying fungi within wood, 59; and HOWARD, N. O., Chemical injuries to white pines, 59; Notes on chemical injuries to the eastern white pine (Pinus strobus L.), 362–368; see YORK, H. H.
SNOW, L. M., A new host for the fire blight organism, Bacillus amylovorus, 517–524
Soil-temperature, influence on Helminthosporium-infection of corn, 30; influence on Helminthosporium-infection of wheat 28; tests with Diplodia zeae, 29; under steam-pan, 53
Solanum melongena, leaf spot of, 61–5
Southern Division, American Phytopathological Society, meeting of, see American Phytopathological Society.
Soybean, new host of Bacterium solanacearum, 98
SPAULDING, PERLEY, Notes on Cronartium ribicola, 46; Viability of telia of Cronartium ribicola in early winter, 221–224
Speckled leaf blotch of wheat, 558–584; Speckled blotch of oats, caused by Leptosphaeria, 450–470
Sphaerotheca pannosa, reported from Crimea, 534
Spinach, host of Puccinia subnitens, 446; Fusarium blight of, 447
Spore, content of upper air, 44; germination in rusts, 471–488; staining method for, 492–494
Sporidia, potential production of, in Cronartium ribicola, 298–9
Spray, alkaline, effect on size of sweet cherries, 104; materials, less common, 251
Spraying for celery blight, 48; of potato, cooperative project, 241–8
Staining method, for fungous hyphae, 440–441; for germinating spores, 492–494
STAKMAN, E. C., and AAMODT, O. S., The effect of fertilizers on the development of stem rust of wheat, 31; and HENRY, A. W., CHRISTOPHER, W. N., and CURRAN, G. C., Observations on the spore content of the upper air, 44
Stalk rot, of corn, control by seed-disinfection, 30
Steam pan, soil temperature under, 53
Stem rust resistance to, in oats, 108; inherited resistance to, in wheat, 31, 32
STEVENS, F. L., A fungus destructive to asphalt shingles, 497
STOKDYK, E. A., see BISBY, G. R.
Stomata, relation to potato tipburn, 305–33
Stone-fruits; bacterial gummosis of, 103; cross-inoculation with Cladosporium, 49

STONE, R. E., Leaf scorch or mollisiose of the strawberry, 375–380; and J. E. HOWITT, Experiments with winter blight or streak of tomatoes, 41; see J. E. HOWITT
STOVER, W. G., The relation of soil temperature to the development of the seedling blight of corn caused by Helminthosporium sp., 30
Strawberry, host of nematode, 105; 445; leaf scorch, 375–380; host of Marsonia potentillae, 375–380; host of Mollisia earliana, 375–380
Streak, potato, transmission of, 41; of winter blight of tomato, control of, 41
Sugar-beet, late blight of, 443
Sulfur, inoculated and uninoculated, for potato scab, 38
Sulfuric acid, for angular leaf spot of cotton, 20
Sunflower, rust of, 44
Synchytrium papillatum, on Erodium moschatum, 442

Take-all disease, of cereals, 66–88; of grasses and cereals, 27; of wheat, caused by Ophiobolus cariceti, 27
TAUBENHAUS, J. J., Recent studies of Texas root rot of cotton, 250
TAYLOR, MINNIE, Potential sporidia production per unit in Cronartium ribicola, 298–9
Temperature, as a factor in growth of Diplodia zeae, 29; of soil in development of Helminthosporium in corn, 30; of soil in development of Helminthosporium in wheat, 28; under steam pan, 53; relation to spore germination 353–361; 473–474; of soil, in relation to Flax wilt, 409–413; relation to distribution of Fusarium diseases, 413; of air, relation to mosaic diseases, 438–440; relation to growth of Corticium vagum, 444
TEODORO, N. G., Pathogenicity of Macrosporium parasiticum, 50
Thielavia basicola, on watermelon, 445
Thielaviopsis paradoxa, on cocoanut palm, 398
THOMAS, H. E., see FITZPATRICK, H. M.
Tilletia tritici, infection in wheat, 108, 337–52
Tipburn, potato, caused by Empoasca mali Le Baron, 181–4; in northeastern Maine, 36; relation to water pores and stomata, 305–33
TISDALE, W. B., see WALKER, J. C., see JONES, L. R.
TISDALE, W. H., Recent investigations on rice disease in the United States, 108
Tobacco, effect of dilutions of Bacterium tumefaciens upon, 56; leaf spot of, 99; non-parasitic leaf-spot of, 52; wild-fire disease of, 51
Tomato wilt organism, spread of, 428–434
Torula ligniperda, in wood of various species, 369–374
Tomato, Isaria rot of, 167–72; mosaic, over-wintering of, 41; new Alternaria spot of, in California, 146–8; winter blight or streak, control of, 41
Transit diseases, 105
Triticum vulgare, crosses of, and resistance to black stem rust, 31
Tropaeolum majus L., inoculated to produce fasciation, 265–70
TRUESDELL, H. W., Plant pathology in Crimea, 533–535
Tsuga canadensis, host of Torula ligniperda, 370
Tylenchus dipsaci, a parasitic nematode, 52; on alfalfa, 105; on strawberries, 445

Urediniospores of corn rust, 89–97
Ustilago avenae, factors in spore germination, 45

VAUGHAN, R. E., and JONES, F. R., Colletotrichum pisi Pat. on garden peas, 47
Venturia spp. reported from Crimea, 533
Verticillium sp. relation to peach wilt disease, 56
Vitis quadrangularis, host of Schizonella **colemani**, 435–438

WALKER, J. C., and TISDALE, W. B., Further notes on the occurrence of cabbage black leg, 43

WALKER, M. N., see DOOLITTLE, S. P.
Walnut blight, infection by and resistance to, 106
Wart disease of potato, 38
Water requirement of wheat affected by rust infection, 31
Watermelon, host of Thielavia basicola, 445
Weather, its relation to citrus scab, 57; relation to potato diseases, 445
WEBER, G. F., Studies on corn rust, 89–97; studies on Septoria diseases of cereals and certain grasses, 44; Septoria diseases of cereals, 449–470; Septoria diseases of wheat, 537–584
WEIMER, J. L., see HARTER, L. L.
WEISS, FREEMAN, The effect of rust infection upon the water requirement of wheat, 31; and ORTON, C. R., Progress notes on potato wart disease investigations, 38
WELLES, C. G., Cercospora leaf spot of eggplant, 61–65
Weymouth pine, relation to Cronartium ribicola, 302–4
Wheat, "black point" of, 34; blight of seedlings by Fusarium culmorum, 28; bunt infection of, 108; 337–52; copper carbonate dust treatment of seed for smut, 36; correlated inheritance of rust resistance and winter-spring habit of growth, 32; effect of delayed planting on formalin-treated seed, 173–9; foot-rot disease in Kansas, 27; glume blotch of, 537–558; Helminthosporium disease, 28; 447; inherited resistance to stem rust, 31, 32; rust infection affecting water requirement of, 31; scab organism, relation of growth to hydrogen-ion concentration, 290–4; seed, treated with formalin, germination of, 173–9; Septoria diseases of, 537–584; Septoria nodorum on, 537–558; Septoria tritici on, 558–584; Speckled leaf blotch of, 558–584; Stem rust, effect of fertilizers on, 31; take-all of, caused by Ophiobolus cariceti, 27
WHETZEL, H. H., note of his resignation, and opinion as to present administrative policies, 499–500
White Pine blister rust, conditions necessary for infection of pine, 148–150; effect on currant, 45; viability of telia of, in winter, 221–4; distribution in Pacific Northwest, 448
White pine, see Pinus strobus
Wildfire disease of tobacco, 51
Wilt, bacterial, of cucumber, 143–6; of peach, 56
WINGARD, S. A., A yeast parasitic on lima beans, 47; Yeast-spot of lima beans, 525–532
Winter blight or streak, of tomatoes, 41
WOLF, F. A., Additional hosts for Bacterium solanacearum, 98; A leaf spot disease of tobacco caused by Phyllosticta nicotiana E. and E., 99–101

Xylaria sp. susceptibility of app'e rootstocks to, 54

Yarrow (Achillea millefolium), parasitized by Comandra pallida, 104
Yellow dwarf, of potato, 39, 123–132
"Yellows" of celery, variety resistant to, 48; peach, dissemination of, 140–2; (leafcurl) of raspberry, 253–64
YORK, H. H., and SNELL, W. H., Experiments in the infection of Pinus strobus with Cronartium ribicola, 148–150

Zea mays, fusaria of, 251; Fusarium-infected, tested in improved germinator box, 30; kernel rot of, 25; mosaic in Arkansas, 250–252; parasite of, Diplodia zeae, 29; Pseudomonas dissolvens on, 497–499; root and stalk rot, control of by seed-disinfection, 30; rust, studies on, 89–97; seedling blight caused by Helminthosporium sp., 30
ZELLER, S. M., Die-back of loganberry in the Northwest, 104; and OWENS, C. E., European canker in the Pacific Coast States, 105; Mophological differences between Nectria galligena Bres. and N. coccinea (ditissima), 442; A "plum pocket" on Prunus subcordata in Oregon, 443

ERRATA

Page 165, line 1 of legend should read "Fig. 2. *Macrosporium carotae*. A. Stem of a mature carrot plant, etc."
Page 279, line 34, "not likely" should be "unlikely."
Page 281, line 13, "Bordeaux Burgundy mixture" should be "Bordeaux or Burgundy mixture."
Page 381, title heading "Roselinia" should be "Rosellinia"
Page 384, line 14, "Roselinia" should be "Rosellinia."
Page 385, page heading, "Roselinia" should be "Rosellinia."
Page 390, line 15 "sore" should be "spore."
Page 396, line 18, "citullina" should be "citrullina."
Page 397, line 11, "Masse" should be "Massee."
Page 434, line 22, "Edgertown" should be "Edgerton."
Page 499, footnote belongs to bottom of page 497.